

# Towards the discovery of CP violation in the leptonic sector with T2K-II and the Near Detector Upgrade

Director: Claudio Giganti (LPNHE Paris)  
Co-Director: Sara Bolognesi (IRFU/DPhP)

T2K is a long-baseline neutrino oscillation experiment (LBL- $\nu$ ) taking data in Japan [1]. The goal of LBL- $\nu$  experiments is to do precise measurements of neutrino oscillation parameters, that depend on the ratio between the neutrino energy (E) and the distance between the production and the detection of neutrinos (L) by producing a neutrino beam in which both, L and E, are optimized.

To produce a neutrino beam, an accelerator is used to accelerate protons that hit a target leading to the emission of hadrons. These hadrons, mostly pions, are selected in charge by a system of magnetic horns and focused on a decay tunnel where they decay after few tenths of meters. Their decay produces a pure beam of muon neutrinos (or antineutrinos) that is observed at a Near Detector and at a Far Detector at a distance of few hundreds of km. In the case of T2K, the Near Detector is ND280 and it is a multipurpose magnetic detector used to predict the neutrino spectra in absence of oscillations, while the Far Detector is Super-Kamiokande, a Water Cherenkov detector at a distance of 295 km from the neutrino production point.

T2K was the first experiment to detect neutrino oscillations in appearance mode by observing the appearance of electron neutrinos [2, 3] and it is currently searching for CP violation in the leptonic sector by comparing appearance probabilities for electron neutrinos and electron antineutrinos. In 2020 T2K reported first hints of CP violation in the leptonic sector [4]. This measurement, if confirmed by more data, could help explaining one of the deepest mysteries in our understanding of the Universe, the matter-antimatter asymmetry. CP violation in the leptonic sector can only be discovered with long-baseline experiments and it is the goal T2K of and of its successor, Hyper-Kamiokande (HK) [5], a detector that will be 10 times larger than SK. HK is under construction in Japan and will start taking data in 2027.

In order to measure CP violation effects, an exceptional understanding of systematic uncertainties is needed. In particular, a careful modeling of the interactions between neutrinos and the nuclei (neutrino cross-sections) and of the differences between neutrinos and antineutrinos is required.

To scrutinize the hints of CP violation reported in 2020, T2K is finalizing the upgrade of the Near Detector that will lead to the second phase of the experiment, T2K-II. The ND280 Upgrade [6] consists in a new active target (Super-FGD) and two new Time Projection Chambers (HA-TPC) instrumented with Resistive MicroMegas modules (ERAM) that will be installed in J-PARC by the end of 2022 and will start collecting neutrino data in 2023. The analyses of these data is the main goal of the thesis.

The ND280 Upgrade will have three main advantages with respect to the current ND280:

- Better efficiency to reconstruct muons emitted at high angle with respect to the neutrinos.
- Better efficiency to reconstruct the hadronic part, reducing the threshold for the reconstruction of protons and pions as well as the capability to reconstruct neutrons from the time of

flight between the neutrino interaction and the neutron re-scattering in the detector [7].

- Better capability to distinguish electron from gammas in order to select a high purity sample of electron neutrino interactions.

The reconstruction of the hadrons will help to better reconstruct the energy of the neutrinos and to improve our knowledge of the nuclear processes on-going when neutrinos interact with nuclei. The capability of selecting a clean sample of electron neutrino and antineutrinos will allow to study differences in the cross-section between electron and muon neutrinos, one of the main systematic uncertainties for future long-baseline experiments. This work will hence be crucial also for Hyper-Kamiokande that will use ND280 as one of its near detectors.

## **Thesis directors, topics and expected timeline**

The supervisors of this thesis are the coordinator of the ND280 Upgrade (Claudio Giganti, LPNHE) and the T2K analysis coordinator (Sara Bolognesi, IRFU/DPhP) thus ensuring an excellent supervision for both, hardware and analysis related topics.

The T2K LPNHE and IRFU/DPhP groups collaborate since the beginning of T2K. Our groups are leading the sensitivity studies and the preparation of the analyses tools for the Upgrade [8], as well as the production of the ERAM modules and of the HA-TPC electronics, the development of the HA-TPC reconstruction tools and the analysis of test beam data [9, 10].

While the thesis is mostly devoted to the analysis of the ND280 Upgrade data, it is expected that the selected PhD candidate will also contribute to the installation and the commissioning of the ND280 Upgrade in Japan during the first year of the thesis, so an interest on both, analysis and hardware activities is suitable.

By Summer 2023 we expect to collect enough statistics with the ND280 Upgrade to include first inputs to the T2K Oscillation Analyses. The analyses topics include exploiting the full reconstruction of the final state of Quasi-Elastic muon neutrino or antineutrino interactions (muons plus protons or neutrons) to reduce systematic uncertainties in T2K oscillation analysis and work on the selection electron neutrino interactions to constraint differences between electron and muon neutrino interactions that constitute one of the main systematics in the search for CPV in T2K-II and in future long-baseline experiments.

An additional topic will be the combination of muon and electron neutrino selected to perform an oscillation analysis at ND280 searching for sterile neutrinos. We published a first search of sterile neutrinos at ND280 in [11] and we expect that the upgraded ND280 will have several advantages, including a larger target mass, a lower threshold to reconstruct leptons, better performances in distinguishing electrons from gammas, and the larger exposure that will be collected in T2K-II.

In summary, we should stress that the timing of the proposed PhD thesis matches perfectly with the ND280 Upgrade schedule and provides a possibility to make a significant and highly visible contribution to the T2K-II experiment.

## References

- [1] K. Abe et al. “The T2K Experiment”. In: *Nucl. Instrum. Meth. A* 659 (2011), pp. 106–135. DOI: 10.1016/j.nima.2011.06.067. arXiv: 1106.1238 [physics.ins-det].
- [2] K. Abe et al. “Indication of Electron Neutrino Appearance from an Accelerator-produced Off-axis Muon Neutrino Beam”. In: *Phys. Rev. Lett.* 107 (2011), p. 041801. DOI: 10.1103/PhysRevLett.107.041801. arXiv: 1106.2822 [hep-ex].
- [3] K. Abe et al. “Observation of Electron Neutrino Appearance in a Muon Neutrino Beam”. In: *Phys. Rev. Lett.* 112 (2014), p. 061802. DOI: 10.1103/PhysRevLett.112.061802. arXiv: 1311.4750 [hep-ex].
- [4] K. Abe et al. “Constraint on the matter–antimatter symmetry-violating phase in neutrino oscillations”. In: *Nature* 580.7803 (2020). [Erratum: *Nature* 583, E16 (2020)], pp. 339–344. DOI: 10.1038/s41586-020-2177-0. arXiv: 1910.03887 [hep-ex].
- [5] K. Abe et al. “Hyper-Kamiokande Design Report”. In: (May 2018). arXiv: 1805.04163 [physics.ins-det].
- [6] K. Abe et al. “T2K ND280 Upgrade - Technical Design Report”. In: (Jan. 2019). arXiv: 1901.03750 [physics.ins-det].
- [7] L. Munteanu et al. “New method for an improved antineutrino energy reconstruction with charged-current interactions in next-generation detectors”. In: *Phys. Rev. D* 101.9 (2020), p. 092003. DOI: 10.1103/PhysRevD.101.092003. arXiv: 1912.01511 [physics.ins-det].
- [8] S. Dolan et al. “Sensitivity of the Upgraded T2K Near Detector to constrain neutrino and anti-neutrino interactions with no mesons in the final state by exploiting nucleon-lepton correlations”. In: (Aug. 2021). arXiv: 2108.11779 [hep-ex].
- [9] D. Attié et al. “Performances of a resistive Micromegas module for the Time Projection Chambers of the T2K Near Detector upgrade”. In: *Nucl. Instrum. Meth. A* 957 (2020), p. 163286. DOI: 10.1016/j.nima.2019.163286. arXiv: 1907.07060 [physics.ins-det].
- [10] D. Attié et al. “Characterization of resistive Micromegas detectors for the upgrade of the T2K Near Detector Time Projection Chambers”. In: *Nucl. Instrum. Meth. A* 1025 (2022), p. 166109. DOI: 10.1016/j.nima.2021.166109. arXiv: 2106.12634 [physics.ins-det].
- [11] K. Abe et al. “Search for short baseline  $\nu_e$  disappearance with the T2K near detector”. In: *Phys. Rev. D* 91 (2015), p. 051102. DOI: 10.1103/PhysRevD.91.051102. arXiv: 1410.8811 [hep-ex].